



Trillium  
Gift of Life  
Network

---

# Ontario's Adult Referral and Listing Criteria for Small Bowel and Liver-Small Bowel Transplantation

---

Version 3.0

---

Trillium Gift of Life Network

---

# Ontario's Adult Referral & Listing Criteria for Small Bowel and Liver-Small Bowel Transplantation

## PATIENT REFERRAL CRITERIA:

The patient referral criteria are criteria for which a Health Care Provider would utilize to refer a patient to a transplant centre for assessment. The criteria identified below are the agreed upon conditions for which an adult patient should be referred for intestine transplant assessment. Patients with intestinal failure and one of the following should be referred for assessment for intestine or liver-intestine transplantation.

- 1) **Hyperbilirubinemia:** Referral for intestine transplant assessment should be considered for patients with persistent hyperbilirubinemia (>80 mmol/l) for a period of more than 2 weeks that is not associated with sepsis.
- 2) **Thrombosis:** Patients with thrombosis of 2 of 4 upper body central veins or extensive mesenteric venous thrombosis with secondary liver or intestine impairment should be referred for transplant assessment.
- 3) **Resection:** Referral for small bowel transplant assessment should be considered for patients with massive small bowel resection (>75% of small bowel).
- 4) **Unmanageable Co-morbidities:** Patients with severely diseased bowel and unmanageable co-morbidities including recurrent septic episodes or intractable severe abdominal fistulae or unmanageable fluid and electrolyte abnormalities should be referred for transplant assessment.
- 5) **Diagnostic Uncertainty:** Patients with continuing prognostic or diagnostic uncertainty should be referred for transplant assessment.
- 6) **Quality of life:** Patients with extremely poor quality of life should be referred for transplant assessment.
- 7) **Extensive Local Tumours:** Patients with otherwise unresectable tumors with no or low grade malignant potential (desmoids tumors, lymphangiomatosis) should be referred for transplant assessment.
- 8) **Consent:** Patients or family members may request a referral for transplant assessment.

## PATIENT LISTING INDICATIONS:

Each patient is assessed individually for their suitability for transplantation by the transplant program. The criteria identified below are the agreed upon conditions for which a patient with intestinal failure may be eligible to be waitlisted for intestine transplantation in Ontario. Not all referred patients will be listed for transplantation.

- 1) **Progressive liver disease:** Listing for transplant may be considered for patients with progressive intestinal failure associated liver disease with plasma bilirubin >100mmol/l and signs of portal hypertension, or synthetic liver dysfunction with coagulopathy.
- 2) **Sepsis:** Patients with recurrent life threatening episodes of sepsis resulting in multi-organ failure, metastatic infectious foci, or acquisition of flora with limited antibiotic sensitivities, are eligible to be waitlisted for transplantation.
- 3) **Vascular Access:** Listing for transplant may be considered for patients with loss of more than 50% of the standard central venous access sites.
- 4) **Surgical Disasters:** Patients with intractable, severe, abdominal fistulae with poor quality of life or patients with extensive mesenteric venous thrombosis may be eligible to be waitlisted for transplantation.
- 5) **Extensive Local Tumours:** Patients with otherwise unresectable tumors with no or low grade malignant potential (desmoids tumors, lymphangiomas) are eligible to be waitlisted for transplantation.
- 6) **Quality of life:** Listing for transplant may also be considered for patients with intestinal failure with high morbidity and extremely poor quality of life. Listing based on this indication is appropriate for a carefully selected group of patients.

## PATIENT LISTING CONTRAINDICATIONS:

The contraindications for transplantation identified below are the agreed upon criteria for which a patient would not be eligible to be waitlisted for intestine transplantation in Ontario.

- 1) **Co-Morbidities:** Patients with any co-morbidity that decreases the likelihood of surviving 5 years post-transplant to below 50% or whom which the peri-operative risk is deemed unacceptably high, are not candidates for intestine transplantation.
- 2) **Consent:** Patients who do not want a transplant should not be listed for intestine transplantation.
- 3) **Post-Transplant Care:** Patients with an inadequate or unsafe post-transplant care plan are not candidates for intestine transplantation.
- 4) **Psychosocial Considerations:** Patients who display any of the following are not candidates for intestine transplantation:
  - Unstable psychiatric disorder, especially one likely to interfere with compliance;
  - Any alcohol and/or illicit drug misuse within six months;

- For patients with alcohol associated liver disease: inability to absolutely abstain from alcohol and/or illicit drug use for six months
- Previous documentation or current unwillingness or inability to follow the advice of health professionals; Intestine transplantation should be delayed until patients have demonstrated adherence to therapy for at least 6 months
- Social support/compliance issues prohibiting adherence to post-operative medications and/or follow-up care.

5) **Malignancy:** Patients with active malignancy are not candidates for transplantation. In general, patients with a previous history of malignancy must be tumour free before proceeding with transplantation. Most transplant candidates should wait a period of time between successful treatment and transplantation.

6) **Pulmonary Disease:** Patients with the following respiratory conditions and severity are not candidates for transplantation:

- Requirement for home oxygen therapy (Grade C);
- Uncontrolled asthma (Grade C);
- Severe cor pulmonale; or,
- Severe chronic obstructive pulmonary disease (COPD), pulmonary fibrosis or restrictive disease with any of the following parameters:
  - Best forced expiratory volume in 1 s (FEV<sub>1</sub>) < 25% predicted value;
  - PO<sub>2</sub> room air < 60 mmHg with exercise desaturation, SaO<sub>2</sub> < 90%;
  - > 4 lower respiratory infections in the last 12 months;
  - Moderate disease with evidence of progression.

7) **Cardiac Disease:** Patients with inoperable valve disease should not be considered for transplant. Patients with severe irreversible (non-uremic) cardiac dysfunction should not be listed for intestinal transplantation alone.

All patients should be assessed for the presence of ischemic heart disease (IHD) before intestine transplantation. The minimum required investigations include history, physical examination, electrocardiogram (ECG) and a chest radiograph. Intestine transplantation is contraindicated in patients with IHD in the following situations:

- Patients with progressive symptoms of angina;
- Patients with a myocardial infarction within 6 months;
- Patients without an appropriate cardiac workup;
- Patients with severe diffuse disease, especially with positive non-invasive tests in whom intervention is not possible and in whom expected survival is sufficiently compromised so that transplantation is not reasonable.

8) **Peripheral Vascular Disease:** Patients with large uncorrectable abdominal aneurysms, severe occlusive common iliac disease, active gangrene or recent atheroembolic events are not candidates for intestine transplantation.

**Version Control:**

<b>Name of Document</b>	Ontario’s Adult Referral and Listing Criteria for Small Bowel and Liver-Small Bowel Transplantation
<b>Version 1.0</b>	Created November 2012
<b>Version 2.0</b>	Updated July 2015
<b>Version 3.0</b>	Updated November 2017
<b>Recommended Next Review</b>	November 2019
<b>Approved By</b>	Provincial Liver and Small Bowel Working Group Transplant Steering Committee